CLAIMS

What is claimed is:

1. A compound of the formula

5

$$\begin{array}{c|c} X & \stackrel{L}{\downarrow} \\ X & \stackrel{L}{\downarrow} \\ X & \stackrel{L}{\downarrow} \\ L^{1} & R \end{array}$$

wherein:

10

M is selected from the group consisting of Os and Ru;

R¹ is hydrogen;

Sub

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

15

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

20

2. A compound according to claim 1, wherein the substituted alkyl includes one or more functional groups selected from the group consisting of aryl, alcohol, thiol, ketone, aldehyde,

ester, ether, amine, imine, amide, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, and halogen.

5

3. A compound according to claim 1, wherein the substituted aryl includes one or more functional groups selected from the group consisting of alkyl, aryl, alcohol, thiol, ketone, aldehyde, ester, ether, amine, imine, amide, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, and halogen.


10 Juh

4. A compound according to claim 1 wherein R is selected from the group consisting of

(a) hydrogen;

15

- (b) C₁-C₂₀ alkyl;
- (c) aryl;
- (d) C_1 - C_{20} alkyl substituted with one or more groups selected from the group consisting of aryl, halide, hydroxy, C_1 - C_{20} alkoxy, and C_2 - C_{20} alkoxycarbonyl; and (e) aryl substituted with one or more groups selected from the group consisting of C_1 - C_{20} alkyl, aryl, hydroxyl, C_1 - C_5 alkoxy, amino, nitro, and halide.



5

- 6. A compound according to claim 5, wherein R is phenyl.
- 7. A compound according to claim 4, wherein R is selected from the group consisting of hydrogen, methyl, ethyl, n-butyl, iso-propyl, -CH₂Cl, -CH₂CH₂CH₂OH, and -CH₂OAc.

Subject of the subjec

15

- 8. A compound according to claim 1, wherein L and L¹ are independently selected from the group consisting of phosphine, sulfonated phosphine, phosphite, phosphinite, phosphinite, arsine, stibine, ether, amine, amide, sulfoxide, carboxyl, nitrosyl, pyridine, and thioether.
- 9. A compound according to claim 8, wherein \(\) and \(\mathbb{L}^1 \) are phosphines independently selected from PR³R⁴R⁵ wherein R³ is selected from the group consisting of secondary alkyl and cycloalkyl and wherein R⁴ and R⁵ are independently selected

from the group consisting of aryl, C₁-C₁₀ primary alkyl, secondary alkyl, and cycloalkyl.

10. A compound according to claim 9, wherein L and L¹ are independently selected from the group consisting of - P(cyclohexyl)₃, -P(cyclopentyl)₃, and -P(isopropyl)₃.

11. A compound according to claim 8, wherein L and L¹ are both -P(phenyl)₃.

10 Sub Bont

5

12. A compound according to claim\8, wherein L and L¹ are the same.

13. A compound according to claim 1, wherein X and X¹ are independently selected from the group consisting of halogen, hydrogen; C_1 - C_{20} alkyl, aryl, C_1 - C_{20} alkoxide, aryloxide, C_3 - C_{20} alkyldiketonate, aryldiketonate, C_1 - C_{20} carboxylate, aryl or C_1 - C_{20} alkylsulfonate, C_1 - C_{20} alkylsulfonyl, or C_1 - C_{20} alkylsulfinyl; each optionally substituted with C_1 - C_5 alkoxy or with a phenyl group optionally substituted with halogen, C_1 - C_5 alkoxy;

20

14. A compound according to claim 13, wherein X and X¹ are independently selected from CI, Br, I, H; benzoate, C_1 - C_5 carboxylate, C_1 - C_5 alkyl, phenoxy, C_1 - C_5 alkoxy, C_1 - C_5 alkylthio, aryl, or C_1 - C_5 alkyl sulfonate; each optionally substituted with C_1 - C_5 alkyl or a phenyl group optionally substituted with halogen, C_1 - C_5 alkyl or C_1 - C_5 alkoxy.

15. A compound according to claim 14, wherein X and X¹ are independently selected from the group consisting of CI, CF₃CO₂, CH₃CO₂, CFH₂CO₂, (CH₃)₃CO, (CF₃)₂(CH₃)CO, (CF₃)(CH₃)₂CO, PhO, MeO, EtO, tosylate, mesylate, and trifluoromethanesulfonate.

16. A compound according to claim 15 wherein X and X¹ are both Cl.

17. A compound of the formula

$$X = C R^{1}$$

$$X = C R$$

wherein:

20

15

5

M is selected from the group consisting of Os and Ru; R¹ is hydrogen;

R is a group selected from the group consisting of

- (a) hydrogen;
- (b) C₁-C₄ alkyl;
- (c) phenyl;
- (d) C_1 - C_4 alkyl substituted with one or more groups selected from the group consisting of halide, hydroxy, and C_2 - C_5 alkoxycarboxyl; and

(e) phenyl substituted with one or more groups selected from the group consisting of C_1 - C_5 alkyl, C_1 - C_5 alkoxy, amino, nitro, and halide;

X and X¹ are independently selected from any anionic ligand; and

L and L^1 are independently phosphines of the formula $PR^3R^4R^5$ wherein R^3 is selected from the group consisting of secondary alkyl and cycloalkyl and wherein R^4 and R^5 are independently selected from aryl, C_1 - C_{10} primary alkyl, secondary alkyl and cycloalkyl.

18. A compound according to claim 17, wherein the substituted phenyl is para-substituted.

5

15

19. A compound according to claim 18, wherein R is phenyl or phenyl substituted with a group selected from the group consisting of chloride, bromide, iodide, fluoride, -NO₂, -NMe₂, methoxy, and methyl.

5

- 20. A compound according to claim 19, wherein R is phenyl.
- 21. A compound according to claim 17, wherein R is selected from the group consisting of hydrogen, methyl, ethyl, n-butyl, iso-propyl, -CH₂Cl, -CH₂CH₂CH₂OH and -CH₂OAc.

0

22. A compound according to claim 17, wherein L and L¹ are independently selected from the group consisting of - P(cyclohexyl)₃, -P(cyclopentyl)₃, and -P(isopropyl)₃.

15

23. A compound according to claim 17, wherein X and X¹ are both Cl.

20

24. A compound according to claim 17, wherein R is phenyl,

M is Ru, X and X¹ are both Cl, and L and L¹ are the same and

are selected from the group consisting of -P(cyclohexyl)₃,
P(cyclopentyl)₃, and -P(isopropyl)₃.

25. A compound of the formula

5 wherein:

M is selected from the group consisting of Os and Ru;

R⁹ and R¹⁰ are independently selected from the group

consisting of hydrogen, substituted or unsubstituted alkyl, and

substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

26. A compound according to claim 25, wherein the substituted alkyl includes one or more functional groups selected from the group consisting of aryl, alcohol, thiol, ketone, aldehyde, ester, ether, amine, mine, amide, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, and halogen.

20

15

27. A compound according to claim 25, wherein the substituted arvl includes one or more functional groups selected from the group consisting of alkyl, aryl, alcohol, thiol, ketone, aldehyde, ester, ether, amine, imine, amide, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, and halogen.

28. A compound according to claim 25, wherein R⁹ and R¹⁰ are independently selected from the group consisting of

- (a) hydrogen;
- (b) C₁-C₂₀ alkyl;
- (c) aryl;
- (d) C_1 - C_{20} alkyl substituted with a group selected from the group consisting of halide, aryl, alkoxy, and aryloxy; and
- (e) aryl substituted with a group selected from the group consisting of halide, alkyl, aryl, alkoxy, and aryloxy.
- 29. A compound according to claim 25, wherein M is Ru, R⁹ and R¹⁰ are hydrogen, X and X¹ are Cl, and L and L¹ are the same and are selected from the group consisting of

10

5

15

P(cyclohexyl)₃, -P(cyclopentyl)₃, -P(isopropyl)₃, and -P(phenyl)₃.

30. A process for polymerizing cyclic olefins comprising the step of contacting a cyclic olefin with a compound of the formula

10 wherein:

M is selected from the group consisting of Os and Ru;

R¹ is hydrogen;


R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

20

15



$$\begin{array}{c|c} X & \stackrel{L}{\downarrow} = C \\ X^1 & \stackrel{L}{\downarrow}_1 = C \\ R & \end{array}$$

in the presence of an acyclic olefin, wherein:

M is selected from the group consisting of Os and Ru; R¹ is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

32. A process for synthesizing a cyclic olefin comprising the step of contacting a diene with a compound of the formula

20

15

5

$$\begin{array}{c|c} x & \downarrow \\ x^1 & \downarrow \\ L^1 & R \end{array}$$

wherein:

M is selected from the group consisting of Os and Ru;

R1 is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

33. A process for synthesizing an unsaturated polymer comprising the step of contacting a diene with a compound of the formula

$$\begin{array}{c} X \stackrel{L}{\underset{L^1}{\bigvee}} = C \stackrel{R^1}{\underset{L^1}{\bigvee}} = C \\ \end{array}$$

- 80 -

wherein:

M is selected from the group consisting of Os and Ru;

5

10


R1 is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

34. A process for synthesizing telechelic polymers by metathesis polymerization comprising contacting a cyclic olefin with a compound of the formula



15

5

10

in the presence of an α, ω -diffunctional defin, wherein:

M is selected from the group consisting of Os and Ru;

R¹ is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

5

35. A process for synthesizing olefins by metathesis comprising contacting an acyclic olefin with a compound of the formula

 $\begin{array}{c|c}
X & \downarrow \\
M = C \\
R^1
\end{array}$

10

wherein:

M is selected from the group consisting of Os and Ru; R¹ is hydrogen;

15

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

20

L and L¹ are independently selected from any neutral electron donor.

5

10

15

36. A process for synthesizing olefins by cross metathesis comprising contacting a first acyclic olefin with a compound of the formula

$$X = C$$

$$X^{1} = C$$

$$R^{1}$$

in the presence of a second acyclic olefin wherein:

M is selected from the group consisting of Os and Ru;

R¹ is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alky, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor.

37. A process for synthesizing a compound of the formula

$$X^{1} \bigcup_{k=1}^{L} C \setminus R^{1}$$

comprising the step of contacting a compound of the formula $(XX^1ML_nL^1_m)_p$ with a diazo compound of the formula $RC(N_2)R^1$, wherein:

M is selected from the group consisting of Os and Ru;

R and R¹ are independently selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand;

L and L¹ are independently selected from any neutral electron donor;

n and m are independently 0-3, provided n+m=3; and p is an integer greater than 0.

38. A process according to claim 36, wherein R1 is hydrogen.

20

15

5

39. A process for synthesizing a compound of the formula

$$X \downarrow L \\ X \downarrow M = C$$

$$L_1 \downarrow R^{11}$$

$$R^{12}$$

comprising the step of contacting a compound of the formula

$$X = \begin{bmatrix} 1 \\ M \end{bmatrix} = C \begin{bmatrix} R^1 \\ R \end{bmatrix}$$

with an olefin of the formula

H C R11

10

15

5

wherein:

M is selected from the group consisting of Os and Ru;

R¹ is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

R¹¹ and R¹² are independently selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor;

5

40. A process for synthesizing a compound of the formula

$$\begin{array}{c|c} X & \stackrel{L}{\downarrow} & \\ X^{1} & \stackrel{L}{\downarrow} & \\ L^{1} & & \\ R^{10} & & \end{array}$$

10

15

20

comprising the step of contacting a compound of the formula $(XX^1ML_nL^1_m)_p$ with an acetylene of the formula R^9CCR^{10} , wherein:

M is selected from the group consisting of Os and Ru;

R⁹ and R¹⁰ are independently selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand; and

L and L¹ are independently selected from any neutral electron donor;

n and m are independently 0-3, provided n+m=3; and p is an integer greater than 0.

41. A process for synthesizing a compound of the formula

$$\begin{array}{c|c}
X & \stackrel{L}{\downarrow} & \\
M = C = C \\
\downarrow & \\
R^{10}
\end{array}$$

comprising the step of contacting a compound of the formula

$$X = C R^{1}$$

$$X^{1} = R$$

with a cumulated olefin of the formula

wherein:

M is selected from the group consisting of Os and Ru;


R¹ is hydrogen;

R is selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

R⁹ and R¹⁰ are independently selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted aryl;

15

5



L and L¹ are independently selected from any neutral electron donor.

42. A process for synthesizing a compound of the formula

$$\begin{array}{c|c}
X & \downarrow \\
M = C \\
X^1 & \downarrow \\
L^2 & R
\end{array}$$

10

15

5

comprising the step of contacting a compound of the formula $(XX^1ML_nL^1_m)_p$ with a diazo compound of the formula $RC(N_2)R^1$ in the presence of a neutral electron donor of the formula L^2 , wherein:

M is selected from the group consisting of Os and Ru;

R and R¹ are independently selected from the group consisting of hydrogen, substituted or unsubstituted alkyl, and

substituted or unsubstituted aryl;

X and X¹ are independently selected from any anionic ligand;

20

L, L¹, and L² are independently selected from any neutral electron donor;

n and m are independently 0-3, provided n+m + 3; and





Docket No. CTCH-1620

p is an integer greater than 0.

